

Spis treści

1. Introduction	7
2. Research objective	9
3. Study areas	10
4. Methods	12
4.1 Characterization of sampling sites	12
4.2 Methods of collecting dung beetles	15
4.3 Choice of sampling sites	17
5. Data analysis methods	19
5.1 Choice of methods for assessing the species diversity of collected beetles	19
5.2 Dominance structure	21
5.3 Analysis of vertical ranges	21
5.4 Analysis of habitat preferences	22
6. Results	23
6.1 Faunistic analysis of collected beetles	23
6.2 Species diversity and dominance structures of dung beetle assemblages in the Polish Carpathians	28
6.2.1 Low Beskids	28
6.2.2 Nowy Sącz Basin (Kotlina Sądecka)	28
6.2.3 Rożnów Foothills (Pogórze Rożnowskie)	28
6.2.4 Beskid Sądecki	35
6.2.5 Western Bieszczady	35
6.2.6 Sanok-Turka Mountains (Góry Sanocko-Turczańskie)	35
6.2.7 Pieniny	35
6.2.8 Żywiec Beskid (Beskid Żywiecki)	35
6.2.9 Ciężkowice Foothills (Pogórze Ciężkowickie)	35
6.2.10 Gorce Mountains	35
6.2.11 Island Beskid (Beskid Wyspowy)	43
6.2.12 Strzyżów Foothills (Pogórze Strzyżowskie)	43
6.2.13 Little Beskids (Beskid Mały)	43
6.2.14 Jasło-Krosno Basin (Kotlina Jasielsko-Krośnieńska)	43
6.2.15 Tatras	43
6.2.16 Gorlice Depression (Obniżenie Gorlickie)	43
6.3 Altitudinal distribution of species in the Polish Carpathians	43
6.4 Habitat preferences of recorded species	50
6.5 Overview of collected species	61
6.5.1. Family: Geotrupidae	61

6.5.2. Family: Scarabaeidae	64
6.5.2.1 Subfamily: Aphodiinae	64
6.5.2.2 Subfamily: Scarabaeinae	94
7. Discussion	105
7.1 Faunistic analysis of collected species	105
7.2 Assessment of species diversity and dominance structures	108
7.3 Vertical distribution of dung beetles in the Polish Carpathians	112
7.4 Habitat preferences of dung beetles in the Polish Carpathians	113
7.5 Endangered species in the Polish Carpathians	114
7.6 Final conclusions	116
8. Conclusions	118
9. References	120
Streszczenie	129